

CLASSIFICATION RESTRICTED

CENTRAL INTELLIGENCE AGENCY

RESTRICTED
INFORMATION REPORT

REPORT

STAT

COUNTRY

USSR

DATE DISTR. 25 May 1948

SUBJECT

Scientific Research

NO. OF PAGES 1

PLACE ACQUIRED

USER

NO. OF ENCLS.
(LISTED BELOW)

DATE OF INFORMATION

1941-46

SUPPLEMENT TO
REPORT NO.

THIS DOCUMENT CONTAINS INFORMATION AFFECTING THE NATIONAL SECURITY OF THE UNITED STATES. IT IS UNLAWFUL TO DISSEMINATE OR TO TRANSMIT BY TELETYPE OR TELETYPE EQUIPMENT, OR TO OTHERWISE COMMUNICATE, IN WHOLE OR IN PART, WITH AN UNAUTHORIZED PERSON THE CONTENTS OF THIS DOCUMENT OR ANY INFORMATION CONTAINED THEREIN, OR TO PUBLISH, REPRODUCE OR DISTRIBUTE IT, DIRECTLY OR INDIRECTLY, IN WHOLE OR IN PART, OR TO COMMUNICATE IT TO AN UNAUTHORIZED PERSON, IN VIOLATION OF THE PROVISIONS OF THE EMBARGO ACT, THE SPY ACT, THE TRADE SECRETS ACT, OR THE NATIONAL SECURITY ACT. IT IS UNLAWFUL TO COMMUNICATE THIS DOCUMENT OR ITS CONTENTS IN WHOLE OR IN PART, OR TO PUBLISH, REPRODUCE OR DISTRIBUTE IT, DIRECTLY OR INDIRECTLY, IN WHOLE OR IN PART, OR TO COMMUNICATE IT TO AN UNAUTHORIZED PERSON, IN VIOLATION OF THE PROVISIONS OF THE EMBARGO ACT, THE SPY ACT, THE TRADE SECRETS ACT, OR THE NATIONAL SECURITY ACT.

THIS IS UNEVALUATED INFORMATION FOR THE RESEARCH
USE OF TRAINED INTELLIGENCE ANALYSTS

SOURCE Documentary as indicated. (Information specifically requested.)

RECENTLY PUBLISHED RESEARCH OF THE
CENTRAL PEDIATRIC RESEARCH INSTITUTE,
MINISTRY OF PUBLIC HEALTH, USSR

"Lysozyme of the Saliva during Ontogenesis," I. A. Arshavskiy, E. V. Morachevskiy, S. M. Sutanov, Cent Pediatric Res Inst

"Byull Eksp Biol i Med" Vol 21, No 4, 1946, pp 69-71

Saliva of newborn puppies or of fetuses removed 2-3 days prematurely lacks lysozyme and if cow milk is substituted for mother's milk, lysozyme does not appear in the saliva until the dogs are 2-2½ months old. In suckled puppies considerable quantities of lysozyme are present within 2-3 days and adult levels are reached on the 15th day. This is considered an example of passive immunization.

"Changes of the Total Amino Acids, Tyrosine, and Mercapto Groups in Erythrocytes in Experimental Anemia," T. V. Fokina, Cent Pediatric Res Inst

"Byull Eksp Biol i Med" Vol 11, 1941, pp 449-52

In the blue-blotting type of anemia, the concentration of amino acids fell, and the concentration of SH and tyrosine rose above normal. In phenylhydrazine anemia the concentration of amino acids fell, the SH concentration rose for 3 days, then returned to normal, while the concentration of tyrosine increased for 2-3 days and then decreased in 3-9 days. The results were not completely consistent.

- END -

- 1 -

CLASSIFICATION RESTRICTED

STATE	X NAVY	X NSN&	DISTRIBUTION					
ARMY	X AIR	X						

RESTRICTED